



# Chicago Respiratory Virus Weekly Surveillance Report



Brandon Johnson, Mayor

March 14, 2025

Olusimbo Ige, MD, MS, MPH, Commissioner


*This report summarizes key respiratory virus surveillance indicators. The indicators are compiled from laboratory-based data as well emergency department visit data. All data are preliminary and may change as additional reports are received. Historical and seasonal summary reports can be found here: [Chicago Influenza and Respiratory Virus Surveillance Report](#).*

**A new Respiratory Illness Dashboard is now available on the [CDPH website](#).** This dashboard summarizes information about respiratory virus disease activity in Chicago, with a focus on COVID-19 (caused by the SARS-CoV-2 virus), influenza, and respiratory syncytial virus (RSV). The dashboard will be updated every Friday at 1pm.

## Weekly Surveillance Key Points

- Overall acute respiratory illness activity level in Chicago has decreased from Moderate to Low. Flu activity remains Moderate, and all influenza clinical surveillance indicators continue to decline. COVID-19 and RSV activity remain Low.
- ED visits, hospitalizations, and ICU admissions attributable to the flu continue to decrease from the late January peak. Overall ED visits, hospitalizations, and ICU admissions for RSV continue to decrease. ED visits and hospitalizations for COVID-19 have decreased or remained the same over the past week across all age groups, and ICU admissions have continued to decrease over the last three weeks.
- The percentage of specimens that tested positive for flu remains elevated (11%) but has decreased from a peak of 23% in January. Detections of flu B have continued to increase and represented 35% of all reported flu detections last week. Among flu A that were subtyped for the week 82% were H1N1. Test positivity for RSV has been decreasing since mid-December and continues to decrease. Test positivity for COVID-19 decreased from the previous week.
- Compared to baseline concentrations, the overall wastewater concentration for flu A remains high, flu B remains low, COVID-19 has decreased from high to low, and RSV has increased from moderate to high. It is possible that even as clinical surveillance indicators (e.g., ED visits and hospitalizations) decrease, wastewater data may fluctuate for longer and decrease over an extended period of time because of the persistence of viral shedding from infected persons and/or in wastewater systems.
- The CDC has issued interim estimates of [2024-25 Seasonal Influenza Vaccine Effectiveness](#) (VE) and [2024-25 COVID-19 VE](#), which suggest that this season's influenza and COVID-19 vaccines were effective in preventing medically-attended illness, including hospitalization, in the US. It's not too late to get vaccinated!
- **Flu and COVID-19 vaccine coverage across the city remain low.** See CDPH's [Mid-Season Seasonal Respiratory Vaccine Coverage Report](#) for more details.
- For information on the national bird flu outbreak and local guidance see: the [CDPH H5N1 Bird Flu](#) webpage and [CDPH HAN: Flu Page](#) for clinicians.

**Influenza-Associated ICU Hospitalizations -** In Illinois, *influenza-associated ICU hospitalizations are reportable* as soon as possible but within three days. The graph below shows the weekly number of reported ICU hospitalizations for Chicago residents for previous two seasons. The table summarizes selected characteristics of reported cases for the current week and cumulative for the season.




Group	Week Ending March 8, 2025		Since September 29, 2024	
	#	%	#	%
<b>Citywide</b>	13	100	491	100
<b>Age</b>				
0-4	0	0.0	32	6.5
5-17	2	15.4	33	6.7
18-24	0	0.0	16	3.3
25-49	3	23.1	85	17.3
50-64	0	0.0	100	20.4
65+	8	61.5	225	45.8
<b>Gender</b>				
Male	7	53.8	238	48.5
Female	6	46.0	253	51.5
<b>Race-Ethnicity</b>				
White Non-Latinx	4	30.8	117	23.8
Black Non-Latinx	6	46.2	229	46.6
Latinx	2	15.4	105	21.4
Asian Non-Latinx	1	7.7	21	4.3
Other Non-Latinx	0	0.0	12	2.4
Unknown-Race	0	0.0	7	1.4

**Respiratory Virus Laboratory Surveillance - Current Week and Cumulative** The table below includes respiratory viral PCR tests performed by several hospital laboratories in Chicago as well as two commercial laboratories serving Chicago facilities. Reporting facilities represent nearly half of all acute care hospitals in the city. Data reported include Chicago and non-Chicago residents.

Respiratory Pathogen	Week Ending March 8, 2025		Since September 29, 2024	
	# Tested	% Positive	# Tested	% Positive
Influenza*	3,285	11.2	87,446	11.6
RSV*	2,645	2.8	71,012	6.6
SARS-CoV-2*	2,666	4.2	71,510	4.1
Parainfluenza	1,707	0.7	47,974	1.3
Rhinovirus/Enterovirus	828	9.1	26,078	12.9
Adenovirus	828	1.3	26,050	1.7
Human Metapneumovirus	828	2.8	26,352	0.7
Seasonal Coronaviruses <sup>†</sup>	1,707	3.0	38,354	2.6


\*Represents both dualplex and multiplex PCR data. All other data represents only multiplex panels that include the specified pathogens;† Four seasonal coronavirus strains include 229E, NL63, OC43, and HKU1.

Weekly number of specimens testing positive for influenza by type and subtype (graph) and the number of positive specimens by type and subtype for the current week and cumulative for the season (table).




Type / Subtype	Week Ending March 8, 2025		Since September 29, 2024	
	# Positive	%	# Positive	%
<b>Influenza A</b>	239	64.8	9,702	95.7
(H1N1)pdm09	23	9.6	1,103	11.4
H3N2	5	2.1	694	7.2
Subtyping not performed	211	88.3	7,905	81.5
<b>Influenza B</b>	130	35.2	433	4.3
<b>Total Positive</b>	369	100	10,135	100

**Respiratory Virus Laboratory Surveillance - Seasonal Trends** These graphs show seasonal trends of selected respiratory virus testing data presented in the previous table. Typical seasonal periods when activity tends to increase for influenza and RSV are indicated by shaded areas. Elevated test positivity outside of typical seasonal periods suggests atypical activity, and increased clinician awareness and testing may be warranted. Yearly data can also be used to compare the timing and intensity of viral activity, although changes in testing patterns also influence yearly trends, and data should be interpreted in the context of other surveillance indicators.





**Emergency Department Illness Surveillance In Illinois**, all 185 acute-care hospitals report emergency department visit data in near-real time to the Illinois Department of Public Health (IDPH). By tracking symptoms (or chief complaints) of patients in emergency departments, public health can promptly detect unusual levels of illness to determine whether a response is warranted.



Percent of emergency department visits attributed to **influenza-like illness (ILI)** for residents of Chicago zip codes based on chief complaint data.




Percent of emergency department visits attributed to **COVID-19 diagnoses** for residents of Chicago zip codes based on chief complaint data.




Percent of emergency department visits attributed to **respiratory syncytial virus (RSV)** diagnoses for residents of Chicago zip codes based on chief complaint data. Seasonal trends are displayed for children younger than 5 years old who are most impacted by RSV.



**Outpatient Visit Illness Surveillance\*** Several outpatient clinics throughout Chicago participate in CDC's Influenza-like Illness Surveillance Network ([ILINet](#)) by reporting on a weekly basis the total number of outpatient clinic visits, and of those visits, the number with influenza-like illness (ILI). This graph shows the percent of medically-attended outpatient visits attributed to ILI as reported by ILINet facilities in Chicago.



**Weekly Pediatric Admissions** Emergency department visit data includes information on whether the visit resulted in a hospital admission at any time during the course of the clinical encounter. The syndromes or disease associated with the hospitalization are based on chief complaint and discharge diagnosis codes and do not necessarily represent lab-confirmed cases. The chart below represents hospital admissions among children <18 years-old at Chicago hospitals due to acute respiratory illnesses.



### National and State Respiratory Virus Surveillance

The Centers for Disease Control and Prevention’s [FluView](#) report provides national updates and trends related to influenza activity across the United States, and the National Respiratory and Enteric Virus Surveillance System ([NREVSS](#)) is a voluntary laboratory-based system that monitors temporal and geographic circulation patterns of several respiratory viruses in the U.S. The Respiratory Syncytial Virus (RSV) Hospitalization Surveillance Network ([RSV-NET](#)) is a CDC population-based surveillance system that collects data on severe RSV hospitalizations, including those resulting in ICU admission or death, among children and adults. The Respiratory Virus Hospitalization Surveillance Network ([RESP-NET](#)) monitors laboratory-confirmed hospitalizations associated with influenza, COVID-19, and respiratory syncytial virus (RSV) among children and adults. The [Illinois](#) and [Suburban Cook County](#) influenza surveillance reports are also available online. Current and archived issues of the Chicago Influenza and Respiratory Virus Surveillance Report can be found on the CDPH website [Historical and Seasonal Summary Reports](#).