Serious About Getting to Zero? Then We Must Address Methamphetamine Use Among Black Gay Men

Presented by Keith R. Green, PhD, MSW

Executive Director, Chicago Black Gay Men's Caucus Co-Chair, Chicago Methamphetamine Task Force





Disclosure: Presenter has no financial interest to disclose. This continuing education activity is managed by The St. Louis STI/HIV Prevention Training Center and accredited by Missouri State Medical Association (MSMA) in cooperation with the Chicago Department of Public Health.

Disclaimer

- This presentation addresses substance use among Black gay, bisexual and same gender loving men, with a specific focus on methamphetamines.
- The content of this presentation may be triggering for some, particularly those who struggle with substance use and/or have a loved one who is or has struggled with substance use.
- We encourage you to take care of yourself during (and after) this presentation. If professional assistance is needed, please check in with the conference organizers.

The Effects of Meth Use

- Using meth...
 - Increases sexual desire and pleasure
 - Allows for disinhibition
 - Helps alleviate the effects of stigma related to sex with other men
 - Can contribute to condomless sex
 - Can cause mental distress, physical damage and social isolation

What We Know About Meth Use

- Data on methamphetamine use among Black gay, bisexual and same gender loving (GBSGL) men is limited
- Data from a recent NIDA study suggests that meth use increased among all populations by 22% between 2015 and 2019
- Frequent meth use increased by 66% during this same time period
- Additionally, use of meth with cocaine increased by 60% during this time

The Chicago Methamphetamine Task Force

The mission of the Chicago Methamphetamine Task Force (CMTF) is to address the destructive use of methamphetamine among Black gay, bisexual, and other men in the City of Chicago by:

- Identifying and amplifying quality of life needs for individuals who are vulnerable to methamphetamine use, abuse and addiction
- **Stimulating** dialogue among service providers, public health officials, elected officials, researchers and other key stakeholders to identify, design, implement and evaluate effective support and treatment programs grounded in the principles of harm reduction



CMTF Mission (cont.)

The mission of the Chicago Methamphetamine Task Force (CMTF) is to address the destructive use of methamphetamine among Black gay, bisexual, and other men in the City of Chicago by:

- Advocating for the designation of existing and/or the creation of new funding streams for addressing this escalating crisis; and
- Encouraging public health officials and service providers to develop protocols for methamphetamine use screening and data collection.



CMTF CommunityWide Survey




A cross-sectional, community-wide survey of 276 Black GBSGL men in Chicago was conducted to:

- Understand substance use behaviors and experiences with substance use treatment among this population;
- **Identify** needs among those who are vulnerable to substance use;
- Promote dialogue among service providers;
- Advocate for existing and new funding streams for addressing this escalating crisis;
- **Encourage** public health officials and service providers to develop enhanced protocols for substance use screening and data collection



CMTF Survey Demographics



Key Survey Findings

- Survey participants aged 25-34 more frequently report using substances from each licit and illicit substance category within the past three months to make sex easier, better, last longer, or something similar
- Majority of participants take substances
 either with their sexual/romantic partner or
 in a social setting rather than alone
- Slightly more than **30**% of participants express either currently being in treatment; previously having been in treatment; or previously having been in detox for substance use



Key Survey Findings (cont.)

- More Black gay, bisexual, and other men who have used methamphetamine within the past three months report currently experiencing homelessness than those who report not having used methamphetamine within the past three months
- Current methamphetamine users mostly report residing on the South and West sides of the city
- Participants who have used methamphetamine within the past three months more frequently report simultaneously using cannabis, poppers, alcoholic beverages, party drugs, cocaine, or nicotine/tobacco products



Key Survey Findings (cont.)

Participants report methamphetamine as the substance they most frequently use in conjunction with other licit and illicit substances



More than 20% of participants are currently either experiencing homelessness or living in a shelter



60% of participants are currently unemployed



Relationship between meth use and HIV

Figure 1 | Black Gay, Bisexual, Same Gender Loving Men Who Have Reported Using Methamphetamines in the Last 3 Months






Figure 2 | New HIV Incidences in 2021



Don't Die High Social Marketing Campaign





Don't Die High Social Marketing Campaign



Thank you!

Questions?

Contact info: kgreen@cbgmcaucus.org